

Contrasting copper evolution in ω Centauri and the Milky Way

Donatella Romano^{1*} and Francesca Matteucci²

¹INAF – Osservatorio Astronomico di Bologna, Via Ranzani 1, I-40127 Bologna, Italy

²Dipartimento di Astronomia, Università di Trieste, Via Tiepolo 11, I-34131 Trieste, Italy

Accepted 2007 March 29. Received 2007 March 27; in original form 2007 March 5

ABSTRACT

Despite the many studies on stellar nucleosynthesis published so far, the scenario for the production of Cu in stars remains elusive. In particular, it is still debated whether copper originates mostly in massive stars or type Ia supernovae. To answer this question, we compute self-consistent chemical evolution models taking into account the results of updated stellar nucleosynthesis. By contrasting copper evolution in ω Cen and the Milky Way, we end up with a picture where massive stars are the major responsible for the production of Cu in ω Cen as well as the Galactic disc.

Key words: Galaxy: evolution – globular clusters: individual: ω Centauri – nuclear reactions, nucleosynthesis, abundances – stars: abundances.

1 INTRODUCTION

Almost two decades ago, Sneden & Crocker (1988) and Sneden, Gratton & Crocker (1991) discussed the possible mechanisms for the synthesis of Cu in stars in the light of their high-resolution observations (showing a decline in $[\text{Cu}/\text{Fe}]$ with decreasing $[\text{Fe}/\text{H}]$ in Galactic stars) and favoured a secondary origin for Cu through the weak component of the s-process in massive stars.

The s-process has long been thought to contribute to the production of Cu in stars (see, e.g., Burbidge et al. 1957). The main s-process, operating in low-mass asymptotic giant branch (AGB) stars, does not contribute more than 5 per cent to the synthesis of Cu isotopes (Raiteri et al. 1993; Travaglio et al. 2004) and will be neglected hereinafter. The weak s-process in massive stars depends on metallicity. This mechanism, in fact, requires neutron fluxes originating mainly from the reaction $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$, where the large abundance of ^{22}Ne during core He burning derives from the original CNO nuclei transmuted into ^{14}N in the H burning ashes, followed by double α -capture on ^{14}N . The residual ^{22}Ne , left behind at core He exhaustion, is fully consumed in the O-rich zone during convective shell C burning. A small primary yield of Cu, 5 to 10 per cent of its solar abundance depending on the location of the mass cut, derives from explosive nucleosynthesis in the inner regions of core-collapse supernovae (SNe II).

In the early nineties, only 25 per cent of solar Cu was attributed to the weak s-process in massive stars (Raiteri et al. 1993). It was advanced then (Matteucci et al. 1993; Mishenina et al. 2002) that most solar Cu originated in type Ia supernovae (SNe Ia). Current SNe Ia models, however, predict a negligible production of Cu during thermonuclear explosions (Iwamoto et al. 1999; Travaglio, Hillebrandt & Reinecke 2005).

It has been recently claimed that up-to-date, high-resolution

spectroscopic data for solar neighbourhood stars belonging to different Galactic substructures (halo, thick-disc and thin-disc), bulge-like stars, globular clusters (GCs) and dwarf spheroidal galaxies (dSphs) can all be well understood if Cu is mostly a secondary product of massive stars, with a small primary contribution from explosive nucleosynthesis (Bisterzo et al. 2005). While this conclusion simply rests on the comparison of extant observations with present stellar nucleosynthesis (and Cu yields from stars ultimately evolving to SNe II must still be viewed with great care – see, e.g., Heger et al. 2001; Hoffman, Woosley & Weaver 2001; Rauscher et al. 2002), a detailed description of copper evolution in different galactic structures is still missing from the literature.

In this Letter, we try to fill in this gap by analysing the evolution of Cu in the framework of detailed galactic chemical evolution (GCE) models in which Cu originates mostly from (i) massive stars or (ii) SNe Ia. In particular, we contrast the evolution of copper in the Milky Way with that in the anomalous GC ω Cen. We conclude that scenarios where copper is produced primarily in SNe Ia are no longer supported by the observations. This imposes strong constraints on the astrophysical processes responsible for the synthesis of Cu in stars.

2 MODELS VERSUS OBSERVATIONS: IMPLICATIONS FOR STELLAR YIELDS AND CHEMICAL EVOLUTION TIME SCALES

Nowadays, high-resolution spectroscopic data of $[\text{Cu}/\text{Fe}]$ are available for large samples of field Galactic stars (see Fig. 1, left panel). GCs generally follow the trends defined by halo field giants (Simmerer et al. 2003): a flat distribution, $[\text{Cu}/\text{Fe}] = -0.75 \pm 0.2$, for low-metallicity stars up to $[\text{Fe}/\text{H}] \simeq -1.8$ dex, followed by a linear increase with a slope close to 1 in the metallicity range $-1.5 < [\text{Fe}/\text{H}] < -1$ (GC data are not plotted in Fig. 1). At

* E-mail: donatella.romano@oabo.inaf.it




Figure 1. Left panel: $[\text{Cu}/\text{Fe}]$ in Galactic disc and halo stars, as measured by Prochaska et al. (2000; big empty squares), Mishenina et al. (2002; stars), Reddy et al. (2003; small filled squares), Allende Prieto et al. (2004; small filled upside-down triangles), Bihain et al. (2004; big empty triangles) and Reddy et al. (2006; dots) in the solar neighbourhood. Also shown are $[\text{Cu}/\text{Fe}]$ values for the very metal-poor stars BD +17° 3248 (Cowan et al. 2002; big filled square), HD 115444, HD 122563 (Westin et al. 2000; big filled triangles) and the extremely metal-poor giant CS 22892–052 (Snedden et al. 2003; big star) and TP-AGB star CS 30322–023 (Masseron et al. 2006; pentagon). Right panel: $[\text{Cu}/\text{Fe}]$ in the globular cluster ω Cen. Data from Cunha et al. (2002) and Pancino et al. (2002; filled circles) have been homogenized as discussed in Romano et al. (2007). Superimposed on the data are our model predictions for ω Cen (right panel, thin lines) and for the solar vicinity (left panel, thick lines), in cases where Cu originates mostly from SN explosions (Models 1 and 2, Table 1, long- and short-dashed lines, respectively; both type II and type Ia SN explosions are considered). The effect of completely removing the (primary) explosive SN II contribution is also shown as a dotted line (only in a model for the solar vicinity; left panel, Model 4, Table 1). See text for details on different models' prescriptions.

$[\text{Fe}/\text{H}] > -0.8$, $[\text{Cu}/\text{Fe}]$ jumps above the solar value. Then, a 'bending' appears for disc stars with $-0.8 < [\text{Fe}/\text{H}] < 0$ (Reddy et al. 2003), though there is a hint that $[\text{Cu}/\text{Fe}]$ might start increasing again at higher metallicities (Allende Prieto et al. 2004).

The Galactic GC ω Cen stands as a notable exception. The $[\text{Cu}/\text{Fe}]$ ratios of its most metal-rich stars are definitely lower than the Galactic trend (Cunha et al. 2002; Pancino et al. 2002; see also Figs. 1 and 2), which can be understood as a shift in the $[\text{Cu}/\text{Fe}]$ relation to higher $[\text{Fe}/\text{H}]$. The only other systems known to have unusually low $[\text{Cu}/\text{Fe}]$ ratios are the Sagittarius dSph (McWilliam & Smecker-Hane 2005) and the Large Magellanic Cloud (LMC; Pompéia, Hill & Spite 2005). In particular, the similarity between the Cu values of ω Cen and Sagittarius adds to the list of common chemical peculiarities (spread in $[\text{Fe}/\text{H}]$ values, strong enhancement of s-process elements) which suggest that the two systems followed similar chemical enrichment histories, thus supporting the idea that ω Cen is the surviving nucleus of an accreted dwarf galaxy (McWilliam & Smecker-Hane 2005).

Superposed on the data in Figs. 1 and 2 are different model predictions for ω Cen (thin lines) and the Milky Way (thick lines). These latter refer to the solar neighbourhood region which, in the framework of the adopted model, is represented by a ring 2 kpc wide centred in the Sun. The model of chemical evolution for the Galaxy is the *two-infall model* of Chiappini, Matteucci & Gratton (1997) and Chiappini, Matteucci & Romano (2001) – where details

about model assumptions and basic equations can be found, except for the adopted stellar lifetimes (which are now taken from Schaller et al. 1992) and stellar yields (see the detailed discussion below). The model of chemical evolution for ω Cen is the one described in Romano et al. (2007), assuming that ω Cen is the remnant of an ancient nucleated dSph evolved in isolation and then accreted by the Milky Way. We adopt this simple model even if a recent study by Villanova et al. (2007) suggests that the chemical evolution of ω Cen might be more complex, with a not unique age-metallicity relation. In fact, accounting for the observed spread needs a full dynamical treatment, while the main conclusions on Cu nucleosynthesis in stars are likely to remain the same.

As regards the adopted stellar nucleosynthesis:

- (i) the s-process yields in massive stars vary as functions of both the initial mass and metallicity of the stars; we follow the prescriptions of either Matteucci et al. (1993; Models 1, 2 and 4, Table 1; long-dashed, short-dashed and dotted lines in Fig. 1, respectively) or Nomoto et al. (2007; Model 3, Table 1; solid lines in Fig. 2);
- (ii) explosive yields of Cu of individual core-collapse SNe are taken from either Matteucci et al. (1993, their model M, $M_{\text{Cu SNeII}}^{\text{ej}} = 7 \times 10^{-6} M_{\odot}$, independent of mass and metallicity; Models 1 and 2, Table 1; long- and short-dashed lines in Fig. 1, respectively) or Nomoto et al. (2007; Model 3, Table 1; solid lines in Fig. 2);
- (iii) the amount of Cu in the ejecta of SNe Ia is of $2 \times 10^{-4} M_{\odot}$




Figure 2. Same data as Fig. 1. The $[\text{Cu}/\text{Fe}]$ versus $[\text{Fe}/\text{H}]$ relation in ω Cen is now compared to that for solar neighbourhood stars. Model predictions for ω Cen (thin solid line) and the solar vicinity (thick solid line) now refer to the evolutive picture where most of solar Cu comes from the weak s-process in massive stars (stellar nucleosynthesis prescriptions as in Model 3, Table 1). See text for details.

Table 1. Prescriptions on stellar Cu production adopted by different GCE models.

Model	$M_{\text{Cu weak s}}^{\text{ej}}$	$M_{\text{Cu SNeII}}^{\text{ej}}$	$M_{\text{Cu SNeIa}}^{\text{ej}}$	Notes
1	Matteucci et al. 1993	Matteucci et al. 1993 (their model M)	Matteucci et al. 1993 (their model M)	Long-dashed lines in Fig. 1
2	Matteucci et al. 1993	Matteucci et al. 1993 (their model M)	Iwamoto et al. 1999 (their model W7)	Short-dashed lines in Fig. 1
3	Nomoto et al. 2007	Nomoto et al. 2007	Iwamoto et al. 1999 (their model W7)	Continuous lines in Fig. 2
4	Matteucci et al. 1993	0	Matteucci et al. 1993 (their model M)	Dotted line in Fig. 1

(Matteucci et al. 1993, their model M; Models 1 and 4, Table 1; long-dashed and dotted lines in Fig. 1) or two orders of magnitude lower (Iwamoto et al. 1999, their model W7; Models 2 and 3, Table 1; short-dashed lines in Fig. 1 and solid lines in Fig. 2), and it is independent of metallicity.

Total yields of copper from massive stars by Matteucci et al. (1993, squares) and Nomoto et al. (2007, dots), comprising both the weak component and the explosive one, are shown and compared one another in Fig. 3, for solar metallicity stars (upper panel) and zero metallicity objects (lower panel). At $Z = 0$, the yields reflect the pure explosive contribution. It is seen that, while the explosive yields adopted by Matteucci et al. (1993) do not vary with the initial mass of the star, those computed by Nomoto et al. (2007) vary steeply as a function of mass, especially for $M_{\text{ini}} > 20 M_{\odot}$.

Models where most of solar Cu comes from SNIa explosions following the prescriptions of Matteucci et al. (1993) match very well the observational data for solar neighbourhood stars (see Fig. 1, left panel, thick long-dashed line), but leave the lower $[\text{Cu}/\text{Fe}]$ ratios in ω Cen totally unexplained (see Fig. 1, right panel, thin long-dashed line). When reducing the contribution from SNeIa according to the results of Iwamoto et al. (1999), a better agreement with the relation observed for ω Cen members is found (except, per-

haps, for the two objects lying at the highest metallicities – Fig. 1, right panel, thin short-dashed line), but the agreement with the solar neighbourhood data at $[\text{Fe}/\text{H}] > -1.0$ is completely destroyed (Fig. 1, left panel, thick short-dashed line). If, instead, most of solar Cu is produced through the weak s-process in massive stars, both the solar neighbourhood and ω Cen relations are satisfactorily reproduced (Fig. 2, thick and thin solid lines). It is worth stressing at this point that, in order to best fit the solar abundance of Cu in the framework of our GCE model, we had to slightly increase (by a factor of 1.5) the yields of Cu by Nomoto et al. (2007) for solar metallicity stars. The applied correction factor, however, is well inside current uncertainties on the weak s-process modelling (see Rauscher et al. 2002 for a discussion on this subject). It is also worth emphasizing how dramatically important the production of Cu from core-collapse SNe is in the very early evolution of the Galactic halo. The SNII contribution sets the level of the ‘plateau’ observed at $[\text{Fe}/\text{H}] < -1.8$. This can be clearly seen from Fig. 1 (left panel), where the thick dotted line (Model 4, Table 1) shows the effect of removing any primary Cu production from explosive nucleosynthesis in massive stars from the Milky Way model. In this case, the ‘plateau’ at low $[\text{Fe}/\text{H}]$ is no longer expected, at variance with the observations, and only the steep rise towards high $[\text{Cu}/\text{Fe}]$




Figure 3. Stellar yields of Cu (in units of solar mass) for solar-metallicity and zero-metallicity massive stars from Matteucci et al. (1993, squares) and Nomoto et al. (2007, dots) as functions of the initial mass of the stars. At solar metallicity (upper panel), the yields comprise a weak s-process component as well as an explosive one, while for $Z = 0$ (lower panel) only the explosive component is present.

at high metallicities can be reproduced. Furthermore, notice that when adopting the yields of Nomoto et al. (2007) rather than those of Matteucci et al. (1993), a lower plateau level is obtained (Fig. 2), because of the lower amount of Cu in the ejecta of low-metallicity SNeII predicted by those authors (see Fig. 3, lower panel).

Now we can draw the following conclusions:

(i) After a short phase in which the primary contribution from explosive nucleosynthesis in core-collapse SN dominates, the evolution of Cu in galaxies of different type is regulated mostly by the weak s-process occurring in massive stars.

(ii) Up-to-date, metallicity-dependent stellar yields of Cu by Nomoto et al. (2007) including both a primary contribution from explosive nucleosynthesis and a secondary contribution from the s-process in massive stars allow us to reproduce very well the observed behaviour of $[\text{Cu}/\text{Fe}]$ versus $[\text{Fe}/\text{H}]$ in the solar vicinity as well as in ω Cen.

(iii) When restricting our analysis to a single environment (the solar vicinity) a degeneracy is found in the solution, in the sense that the decline in $[\text{Cu}/\text{Fe}]$ with decreasing $[\text{Fe}/\text{H}]$ can be ascribed to either the reduced extent of the weak s-process in massive stars at low $[\text{Fe}/\text{H}]$, or the delayed Cu production from SNeIa. Only studying objects which have chemically enriched along different evolutionary paths allows us to break the degeneracy.

Finally, in Fig. 4 we compare the abundances of Cu measured in a sample of bulge-like stars by Pompéia (2003) with our model predictions for the inner Galaxy (bulge-like stars are thought to originate near the Galactic bulge). A substantial contribution to the Galactic Cu production from SNeIa (nucleosynthesis prescriptions




Figure 4. Predictions from Models 1 (long-dashed line), 2 (short-dashed line) and 3 (solid line) computed for the inner Galaxy ($R_G = 4$ kpc) compared to measurements of $[\text{Cu}/\text{Fe}]$ in a sample of bulge-like stars (data from Pompéia 2003).

as in Model 1, Table 1; long-dashed line) is clearly ruled out, while models with a minimum Cu production from SNeIa can well fit the data (Models 2 and 3, short-dashed and solid lines, respectively). We conclude that the study of Cu abundances in *different environments*, from both an observational and a theoretical point of view, is of fundamental importance in order to understand the origin of Cu in the universe.

ACKNOWLEDGMENTS

Thanks are due to the referee, Raffaele Gratton, for his careful reading of the manuscript and constructive comments.

REFERENCES

- Allende Prieto C., Barklem P. S., Lambert D. L., Cunha K., 2004, *A&A*, 420, 183
- Bihain G., Israelian G., Rebolo R., Bonifacio P., Molaro P., 2004, *A&A*, 423, 777
- Bisterzo S., Pompéia L., Gallino R., Pignatari M., Cunha K., Heger A., Smith V., 2005, *Nuc. Phys. A*, 758, 284c
- Burbidge E. M., Burbidge G. R., Fowler W. A., Hoyle F., 1957, *Rev. Mod. Phys.*, 29, 547
- Chiappini C., Matteucci F., Gratton R., 1997, *ApJ*, 477, 765
- Chiappini C., Matteucci F., Romano D., 2001, *ApJ*, 554, 1044
- Cowan J. J., et al., 2002, *ApJ*, 572, 861
- Cunha K., Smith V. V., Suntzeff N. B., Norris J. E., Da Costa G. S., Plez B., 2002, *AJ*, 124, 379
- Heger A., Woosley S. E., Martínez-Pinedo G., Langanke K., 2001, *ApJ*, 560, 307
- Hoffman R. D., Woosley S. E., Weaver T. A., 2001, *ApJ*, 549, 1085

- Iwamoto K., Brachwitz F., Nomoto K., Kishimoto N., Umeda H., Hix W. R., Thielemann F.-K., 1999, *ApJS*, 125, 439
- Masseron T., van Eck S., Famaey B., Goriely S., Plez B., Siess L., Beers T. C., Primas F., Jorissen A., 2006, *A&A*, 455, 1059
- Matteucci F., Raiteri C. M., Busso M., Gallino R., Gratton R., 1993, *A&A*, 272, 421
- McWilliam A., Smecker-Hane T. A., 2005, *ApJ*, 622, L29
- Mishenina T. V., Kovtyukh V. V., Soubiran C., Travaglio C., Busso M., 2002, *A&A*, 396, 189
- Nomoto K., Tominaga N., Umeda H., Kobayashi C., Maeda K., 2007, *Nucl. Phys. A*, 777, 424
- Pancino E., Pasquini L., Hill V., Ferraro F. R., Bellazzini M., 2002, *ApJ*, 568, L101
- Pompéia L., 2003, in Lépine J., Gregorio-Hetem J., eds, *Astrophys. Space Sci. Library* Vol. 299, *Open Issues in Local Star Formation*. Kluwer Academic Publishers, Dordrecht, p. P58
- Pompéia L., Hill V., Spite M., 2005, in Hill V., François P., Primas F., eds, *Proc. IAU Symp. 228, From Lithium to Uranium: Elemental Tracers of Early Cosmic Evolution*. Cambridge Univ. Press, Cambridge, p. 519
- Prochaska J. X., Naumov S. O., Carney B. W., McWilliam A., Wolfe A. M., 2000, *AJ*, 120, 2513
- Raiteri C. M., Gallino R., Busso M., Neuberger D., Käppeler F., 1993, *ApJ*, 419, 207
- Rauscher T., Heger A., Hoffman R. D., Woosley S. E., 2002, *ApJ*, 576, 323
- Reddy B. E., Tomkin J., Lambert D. L., Allende Prieto C., 2003, *MNRAS*, 340, 304
- Reddy B. E., Lambert D. L., Allende Prieto C., 2006, *MNRAS*, 367, 1329
- Romano D., Matteucci F., Tosi M., Pancino E., Bellazzini M., Ferraro F. R., Limongi M., Sollima A., 2007, *MNRAS*, 376, 405
- Schaller G., Schaerer D., Meynet G., Maeder A., 1992, *A&AS*, 96, 269
- Simmerer J., Sneden C., Ivans I. I., Kraft R. P., Shetrone M. D., Smith V. V., 2003, *AJ*, 125, 2018
- Sneden C., Crocker D. A., 1988, *ApJ*, 335, 406
- Sneden C., Gratton R. G., Crocker D. A., 1991, *A&A*, 246, 354
- Sneden C., et al., 2003, *ApJ*, 591, 936
- Travaglio C., Gallino R., Amone E., Cowan J., Jordan F., Sneden C., 2004, *ApJ*, 601, 864
- Travaglio C., Hillebrandt W., Reinecke, M., 2005, *A&A*, 443, 1007
- Villanova S. et al., 2007, *ApJ*, in press (astro-ph/0703208)
- Westin J., Sneden C., Gustafsson B., Cowan J. J., 2000, *ApJ*, 530, 783

This paper has been produced using the Royal Astronomical Society/Blackwell Science $\LaTeX$ style file.